



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

**Genome sequence of *Enterobacter turicensis* strain 610/05 (= LMG 23731),
isolated from fruit powder**

Stephan, Roger ; Grim, C J ; Gopinath, G R ; Mammel, M K ; Sathyamoorthy, V ; Trach, L H ; Chase,
H R ; Fanning, S ; Tall, B D

DOI: <https://doi.org/10.1128/genomeA.00996-13>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-87571>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 3.0 Unported (CC BY 3.0) License.

Originally published at:

Stephan, Roger; Grim, C J; Gopinath, G R; Mammel, M K; Sathyamoorthy, V; Trach, L H; Chase, H R; Fanning, S; Tall, B D (2013). Genome sequence of *Enterobacter turicensis* strain 610/05 (= LMG 23731), isolated from fruit powder. *Genome Announcements*, 1(6):e00996-13.

DOI: <https://doi.org/10.1128/genomeA.00996-13>

Genome Sequence of *Enterobacter turicensis* Strain 610/05 (LMG 23731), Isolated from Fruit Powder

Roger Stephan,^a Christopher J. Grim,^b Gopal R. Gopinath,^b Mark K. Mammel,^b Venugopal Sathyamoorthy,^b Larisa H. Trach,^b Hannah R. Chase,^b Séamus Fanning,^c Ben D. Tall^b

Institute for Food Safety and Hygiene, University of Zurich, Zürich, Switzerland^a; CFSAN, FDA, Laurel, Maryland, USA^b; UCD Centre for Food Safety, School of Public Health, Physiotherapy & Population Science, University College, Dublin, & WHO Collaborating Centre for *Cronobacter*, Belfield, Dublin, Ireland^c

We report the draft genome sequence of *Enterobacter turicensis* strain 610/05 (LMG 23731), isolated from fruit powder. The draft genome has a size of 4,182,790 bp and a G+C% content of 58.0.

Received 24 October 2013 Accepted 31 October 2013 Published 5 December 2013

Citation Stephan R, Grim CJ, Gopinath GR, Mammel MK, Sathyamoorthy V, Trach LH, Chase HR, Fanning S, Tall BD. 2013. Genome sequence of *Enterobacter turicensis* strain 610/05 (LMG 23731), isolated from fruit powder. *Genome Announc.* 1(6):e00996-13. doi:10.1128/genomeA.00996-13.

Copyright © 2013 Stephan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](#).

Address correspondence to Roger Stephan, stephanr@fsafety.uzh.ch.

Stephan et al. (1) reported the isolation from fruit powders of 12 bacterial strains, which were presumptively identified as *Enterobacter sakazakii* (now genus *Cronobacter*) based on colony appearances on *Cronobacter* chromogenic agars. API32E analysis revealed that these isolates were not *Cronobacter* and also that they did not match any existing identification profiles with high confidence. Using a polyphasic analysis scheme, Stephan et al. (1) proposed to classify two of these strains as the novel species *Enterobacter turicensis*. Interestingly, *Cronobacter* comprises a diverse genus of pathogens that cause life-threatening infantile infections, such as brain abscesses, meningitis, necrotizing enterocolitis, and septicemia (2, 3). There is, however, no indication that *E. turicensis* is pathogenic to humans. Not surprisingly, *E. turicensis* was found to share several physiological traits with the genus *Cronobacter*, such as production of a yellow, *Pantoea*-like carotenoid pigment (4) and constitutive metabolism of 5-bromo-4-chloro-3-indolyl-a-D-glucopyranoside, two traits used in the differentiation of presumptive *Cronobacter* colonies.

Recently, Brady et al. (5) proposed that *E. turicensis* be recognized as a new *Cronobacter* species and subsequently, Masood et al. published the draft genome sequence for the type strain of *E. turicensis*, 508/05 (LMG 23730; DSM 18397) (6). Because the taxonomic position of these strains remains unclear, we sequenced *E. turicensis* strain 610/05 (LMG 23731). A library was constructed using a Nextera XT DNA sample preparation kit (Illumina, San Diego, CA), and whole-genome sequencing was performed on a MiSeq benchtop sequencer (Illumina, San Diego, CA), utilizing 500 cycles paired-end version 2 chemistry. FASTQ datasets were trimmed and assembled using default parameters in CLC Genomics Workbench, version 6.0.5 (CLC bio, Aarhus, Denmark). The draft genome of strain 610/05 is 4,182,790 bp, on 262 contigs (>500 bp in size), and has a G+C% content of 58.0. Genomic contigs were annotated using the RAST annotation server (7) to identify RNAs and protein-coding genes. The draft genome of strain 610/05 is predicted to contain 3,857 coding sequences (CDS).

The *E. turicensis* strain 610/05 draft genome sequence was

highly clonal with that of strain LMG 23730^T (average nucleotide identity of 99.98), as reported by Masood et al. (6). The genome possessed a number of noteworthy features, including six chaperone-usher fimbriae; curli, dulcitol, and malonate utilization; and a type III secretion system gene cluster similar in gene content to those of *Pectobacterium carotovorum* and *Pseudomonas syringae*. The genome contained two conjugative plasmids, an IncF (*tra*) plasmid that contains a copper homeostasis operon and an IncN (*virB*) plasmid similar to that found in *Salmonella enteritidis* serovar Agona strain SL483. It should be noted that the IncF plasmid was not homologous to the common virulence plasmid reported among *Cronobacter* spp., which contains a number of genus- and species-specific virulence factors (8).

Nucleotide sequence accession numbers. The whole-genome shotgun project for *E. turicensis* strain 610/05 is available in GenBank under accession number [AXDM00000000](#). The corresponding NCBI Biosample record SAM02319257 is subject to taxonomic revision.

ACKNOWLEDGMENTS

We acknowledge the financial support provided in part through the Irish government's Food Institutional Research Measure (FIRM) grant number 05/R&D/D/363.

REFERENCES

- Stephan R, Van Trappen S, Cleenwerck I, Vancanneyt M, De Vos P, Lehner A. 2007. *Enterobacter turicensis*, sp. nov. and *Enterobacter helveticus* sp. nov. isolated from fruit powder. *Int. J. Syst. Evol. Microbiol.* 57: 820–826.
- Iversen C, Mullane N, McCardell B, Tall BD, Lehner A, Fanning S, Stephan R, Joosten H. 2008. *Cronobacter* gen. nov., a new genus to accommodate the biogroups of *Enterobacter sakazakii*, and proposal of *Cronobacter sakazakii* gen. nov., comb. nov., *Cronobacter malonaticus* sp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov., *Cronobacter* genomospecies 1, and of three subspecies, *C. dublinensis* subsp. *dublinensis* subsp. nov., *Cronobacter dublinensis* subsp. *lausannensis* subsp. nov. and *Cronobacter dublinensis* subsp. *lactaridi* subsp. nov. *Int. J. Syst. Evol. Microbiol.* 58:1442–1447.
- Joseph S, Cetinkayaz E, Drahovska H, Levican A, Figueras MJ, Forsythe SJ. 2012. *Cronobacter condimentii* sp. nov., isolated from spiced meat and

- Cronobacter universalis* sp. nov., a novel species designation for *Cronobacter* sp. genomospecies 1 recovered from a leg infection, water and food ingredients. *Int. J. Syst. Evol. Microbiol.* 62:1277–1283.
4. Lehner A, Grimm M, Rattei T, Ruepp A, Frishman D, Manzardo GG, Stephan R. 2006. Cloning and characterization of *Enterobacter sakazakii* pigment genes and in situ spectroscopic analysis of the pigment. *FEMS Microbiol. Lett.* 265:244–248.
 5. Brady C, Cleenwerck I, Venter S, Coutinho T, De Vos P. 2013. Taxonomic evaluation of the genus *Enterobacter* based on multilocus sequence analysis (MLSA): Proposal to reclassify *E. nimipressuralis* and *E. amnigenus* into *Lelliottia* gen. nov. as *Lelliottia nimipressuralis* comb. nov. and *Lelliottia amnigena* comb. nov., respectively, *E. gergoviae* and *E. pyrinus* into *Pluralibacter* gen. nov. as *Pluralibacter gergoviae* comb. nov. and *Pluralibacter pyrinus* comb. nov., respectively, *E. cowanii*, *E. radicincitans*, *E. oryzae* and *E. arachidis* into *Kosakonia* gen. nov. as *Kosakonia cowanii* comb. nov., *Kosakonia radicincitans* comb. nov., *Kosakonia oryzae* comb. nov. and *Kosakonia arachidis* comb. nov., respectively, and *E. turicensis*, *E. helveticus* and *E. pulveris* into *Cronobacter* as *Cronobacter zurichensis* nom. nov., *Cronobacter helveticus* comb. nov. and *Cronobacter pulveris* comb. nov., respectively, and emended description of the genera *Enterobacter* and *Cronobacter*. *Syst. Appl. Microbiol.* 36:309–319.
 6. Masood N, Moore K, Farbos A, Hariri S, Paszkiewicz K, Dickins B, McNally A, Forsythe S. 2013. Draft genome sequences of three newly identified species in the genus *Cronobacter*, *C. helveticus* LMG23732^T, *C. pulveris* LMG24059, and *C. zurichensis* LMG23730^T. *Genome Announc.* 1(5):e00783–13. doi:10.1128/genomeA.00783-13.
 7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
 8. Franco AA, Hu L, Grim CJ, Gopinath G, Sathyamoorthy V, Jarvis KG, Lee C, Sadowski J, Kim J, Kothary MH, McCardell BA, Tall BD. 2011. Characterization of putative virulence genes on the related RepFIB plasmids harbored by *Cronobacter* spp. *Appl. Environ. Microbiol.* 77:3255–3267.